

IOBC/wprs Bulletin, Vol. 34, 2008:

Working Group "Landscape management for functional biodiversity", Preceedings of the meeting at Bordeaux (France), 14 - 17 May, 2008. Edited by: Walter Rossing. ISBN 978-92-9067-208-1 [vii + 136 pp.].

| | |
|---|-------|
| Preface | iii |
| Contents | v |
| Which biotopes can supply the arable field with natural enemies? <i>Valentina M. Afonina, Wladimir B. Tshernyshev, Olga V. Solovchenko</i> | 1-4 |
| Model for integrating internal and external drivers for dispersal and distribution pattern in carabid beetles <i>Bas Allema, Walter Rossing, Wopke van der Werf, Tibor Bukovinszky, Eveliene Steingröver, Ariena van Bruggen, Joop van Lenteren, Kees Booij</i> | 5-8 |
| Insectary plants to enhance the biological control of <i>Nasonovia ribisnigri</i> and <i>Frank- liniella occidentalis</i> in lettuce <i>Oscar Alomar, Judit Arnó, Rosa Gabarra</i> | 9-12 |
| Hover-Winter: a multi-agent model to simulate the overwintering of a beneficial insect (<i>Episyrphus balteatus</i> , Diptera, Syrphidae) in a heterogeneous landscape <i>Florent Arrignon, Claude Monteil, Jean-Pierre Sarthou, Marc Deconchat, Gérard Balent</i> | 13-16 |
| Gis-based methodology to assess pesticide risk on biodiversity in terrestrial eco- systems <i>Stefania Barmaz, Serenella Sala, Marco Vighi</i> | 17-20 |
| Mapping the ecosystem service of pest control associated with forest in agricultural landscapes; a proof of concept <i>J.M. Baveco, F.J.J.A. Bianchi, W. van der Werf, P.W. Goedhart</i> | 21-23 |
| Vegetation on field margins as a source of predatory mites (Phytoseiidae) for strawberry plantations in Poland: Preliminary observations <i>Zbigniew T. Dabrowski, Danuta Kropczyńska, Jakub Garnis</i> | 25-28 |
| Green bridges over the winter: consequences for <i>Brassica</i> pests <i>Eefje den Belder, Jane Landure, Janneke Elderson, Marian Vlaswinkel, Jeroen Willemse, Frans van Alebeek, Paul van Rijn, Henny van Gorp</i> | 29-32 |
| The impact of agri-environment schemes on cereal aphid control <i>John Holland, Heather Oaten, Steve Moreby, Sue Southway</i> | 33-36 |
| Long-term set-asides and functional biodiversity <i>Erja Huusela-Veistola</i> | 37-40 |
| Botanical and social aspects of conservation headlands in Switzerland <i>Katja Jacot, Lisa Eggenschwiler, Nina Richner, Daniel Schaffner</i> | 41-44 |
| Combining studies on crop mosaic dynamics and pest population dynamics to foster biological control <i>Alexandre Joannon, Aude Vialatte, Chloé Vasseur, Jacques Baudry, Claudine Thenail</i> | 45-48 |
| Explaining participation in multifunctional agricultural activities: an empirical analysis of the Winterswijk region in The Netherlands <i>Roelof A. Jongeneel, Nico B.P. Polman, Louis H.G. Slangen</i> | 49-52 |
| Reservoirs role of some weed plants in the agroecosystem-dominated landscapes of southeastern Europe (Hymenoptera: Braconidae: Aphidiinae) <i>Nickolas G. Kavallieratos, Željko Tomanović, Petr Starý, Christos G. Athassiou</i> | 53-55 |
| Biodiversity of spiders appearing on nettle (<i>Urtica dioica</i> L.) in natural stands <i>Hanna Legutowska, Magdalena Sitko</i> | 57-60 |
| The 'Cake on the plate' syndrome ... and how specialist tansy aphids avoid extinction by hungry predators and parasitoids: insights from the use of polymorphic micro-satellite markers <i>Hugh D. Loxdale, Franklin Nyabuga, Wolfgang W. Weisser</i> | 61-64 |

| | |
|---|---------|
| Determinants for implementing different types of conservation measures in intensive arable regions <i>Juliane Mante, Bärbel Gerowitt</i> | 65-68 |
| Are landscape structures important for the colonization of spinach fields by insects? <i>Rainer Meyhöfer, Thomas Klug, Hans-Michael Poehling</i> | 69-72 |
| Does the spatial density of field margins affect aerially dispersing aphid predators, if so, at what scale? <i>Heather Oaten, John Holland, Barbara Smith, Simon Leather</i> | 73-76 |
| Evaluating predator diversity and abundance in vineyards and the contiguous hedgerows <i>Stefan Otto, Filippo Maria Buzzetti, Giuseppe Zanin, Carlo Duso</i> | 77-80 |
| Biodiversity and its interactions with viticulture in a wine-growing area in the west of France: case study of a local initiative in a controlled origin appellation (AOC Saumur-Champigny) <i>Guillaume Pain, Véronique Beaujouan, Hervé Daniel, David Montembault, Joséphine Pithon, Annie Sigwalt</i> | 81-84 |
| Expression of on-farm functional biodiversity as affected by management and multi-scale agroecosystem disturbance <i>Souzi Roupheal, Anna-Camilla Moonen, Paolo Bàrberi, Ruggero Petacchi, Luigi Boccaccio</i> | 85-88 |
| The effect of source habitats on arable spider communities: is proximity the most important? <i>Ferenc Samu, András Horváth, Éva Szita, Balázs Bernáth, Erika Botos, Kinga Fetykó</i> | 89-92 |
| Stable isotopes as a tool for food web analysis <i>Stefan Scheu</i> | 93-96 |
| Implementation of an agri-environmental scheme providing sown boundary strips in an agricultural landscape of northern Germany <i>Horst-Henning Steinmann, Jan Freese, Sebastian Klimek</i> | 97-100 |
| Some approaches to natural enemies management <i>Wladimir B. Tshernyshev</i> | 101-104 |
| The dynamics of generalist predators in two different six years crop rotation systems: sources and sinks? <i>F. van Alebeek, R. van den Broek, J.H. Kamstra, W. van den Berg, A. Visser</i> .. | 105-108 |
| Ecological infrastructure and polycultures to improve natural control of insect pests in cabbage: first year results <i>Rob van den Broek, Frans van Alebeek, Wim van den Berg</i> | 109-112 |
| Kernel approach for quantifying the spatial extent of the ecosystem service of pest control provided by non-crop habitats in agricultural landscapes <i>W. van der Werf, P. Goedhart, F.J.J.A. Bianchi & H. Baveco</i> | 113-116 |
| Experimenting with landscape management to control pest populations in viticulture <i>Maarten van Helden, Guillaume Pain, Josephine Pithon, Marie-Anne Simonneau</i> | 117-120 |
| Perspectives for functional agro biodiversity in Brussels sprouts <i>Paul van Rijn, Eefje den Belder, Janneke Elderson, Marian Vlaswinkel, Frans van Alebeek</i> | 121-124 |
| Functional agro biodiversity in Dutch arable farming: results of a three year pilot <i>Paul van Rijn, Frans van Alebeek, Eefje den Belder, Felix Wäckers, Jan Buurma, Jeroen Willemse, Henny van Gulp</i> | 125-128 |
| Spatial analysis of greenhouse density in relation to western flower thrips (<i>Frankliniella occidentalis</i>), onion thrips (<i>Trips tabaci</i>) and minute pirate bug (<i>Orius</i> spp.) population in greenhouses <i>Andrea Veres, Ferenc Tóth, Szilvia Orosz, Daniel Kristóf, Kinga Fetykó</i> | 129-132 |
| Tracing food source use by nectarivorous insects <i>Felix L. Wäckers</i> | 133-136 |